



### PRODUCT DATA

#### TYPICAL ANALYSIS:

Protein: 16%

Approximate on a DM basis. Analysis may vary slightly due to grain variances beyond our control.

#### FEEDING RECOMMENDATIONS:

- Introduce Meal Time Alpaca Pellets plus Zinc gradually to allow digestive systems to adjust to the higher starch intake.
- Alpacas should always have access to a quality fibre source such as hay, chaff, pasture or silage to maintain a healthy digestive system.
- Do not allow individual animals to over consume. When feeding a group of animals spread the feed along the trough to allow for easy access by them all.
- Keep feed fresh and ensure feeding containers are kept clean.
- Ensure that Alpacas have access to fresh, clean water at all times.

Suggested Feeding Rates of Meal Time Alpaca Pellets plus Zinc to help prevent Facial Eczema:

- Feed Meal Time Alpaca Pellets plus Zinc from mid-January to mid-April to help with the prevention of Facial Eczema.
- **Do not feed Meal Time Alpaca Pellets plus Zinc for more than 100 days continuously.**

Suggested Feeding Rates to aid in prevention of Facial Eczema:

- Feeding Rate: 3grams/kg of body weight
- Adult Males(80kg): 240grams/head/day
- Adult Females(60kg):180grams/head/day
- Young: 50-150grams/head/day depending on weight
- Cria: 50-100grams/head/day depending on weight

#### INGREDIENTS:

Selected from: Grain and grain by-products, various plant proteins, molasses, minerals, vegetable oils, mold inhibitor, flavors, zinc, vitamins and trace minerals.

#### CONDITIONS OF SALE:

Being used outside their control, this product is sold on the express condition that the producers have no warranty, expressed or implied and are not responsible for damage arising out of its use, whether in accordance with its directions or not.

#### MANUFACTURED BY:

Denver Stock Feeds Ltd, 815 Tremaine Ave, Palmerston North, New Zealand.



# Fresh is Best

## ALPACA + Zinc

PELLETS

Meal Time Alpaca Pellets plus Zinc have been developed as a supplementary feed to help in the prevention of Facial Eczema(FE). These pellets that include zinc should be fed over the most dangerous period for FE from mid-January to mid-April.

FOR ANIMAL TREATMENT ONLY

PLEASE ENSURE PRODUCT IS STORED IN A COOL, DRY AND PEST FREE ENVIRONMENT.

DO NOT FEED TO ANY OTHER ANIMAL SPECIES OTHER THOSE STIPULATED ON THE LABEL.



FOR MORE INFORMATION

FREEPHONE 0800 336 837

VIEW THE FULL MEALTIME RANGE AT

WWW.MEALTIME.CO.NZ

PRODUCED WITH ONLY THE FRESHEST INGREDIENTS